



INTERNATIONAL
CONGRESS 2020
v i r t u a l

Masitinib in severe asthma: Results from a randomized, phase 3 trial

Pascal CHANEZ, MD, PhD.

Aix-Marseille University, Marseille, France

On behalf of the AB07015 Study Group

Conflict of interest disclosure



I have no real or perceived conflicts of interest that relate to this presentation.

I have the following real or perceived conflicts of interest that relate to this presentation:

Affiliation / Financial interest	Commercial company
Grants/research support:	Almirall, Boston Scientific, Boehringer Ingelheim, Centocor, GlaxoSmithKline, AstraZeneca, Novartis, Teva
Honoraria or consultation fees:	Almirall, Boehringer Ingelheim, Johnson & Johnson, GlaxoSmithKline, Merck Sharp & Dohme, AstraZeneca, Novartis, Teva, Chiesi, Sanofi, SNCF, Centocor, Boston Scientific and ALK
Participation in a company sponsored bureau:	Almirall, Boehringer Ingelheim, Johnson & Johnson, GlaxoSmithKline, AstraZeneca, Novartis, Teva, Chiesi, Schering Plough, and Sanofi
Stock shareholder:	
Spouse / partner:	
Other support / potential conflict of interest:	

This event is accredited for CME credits by EBAP and EACCME and speakers are required to disclose their potential conflict of interest. The intent of this disclosure is not to prevent a speaker with a conflict of interest (any significant financial relationship a speaker has with manufacturers or providers of any commercial products or services relevant to the talk) from making a presentation, but rather to provide listeners with information on which they can make their own judgments. It remains for audience members to determine whether the speaker's interests, or relationships may influence the presentation. The ERS does not view the existence of these interests or commitments as necessarily implying bias or decreasing the value of the speaker's presentation. Drug or device advertisement is forbidden.



Masitinib is a first-in-class oral tyrosine kinase inhibitor in severe asthma

- ❖ Masitinib selectively targets mast cell activity (c-Kit, LYN, FYN) and is also a potent inhibitor of PDGFR
- ❖ Strong scientific rationale to target mast cells in severe asthma
- ❖ PDGFR signaling associated with airway remodeling in severe asthma
- ❖ Masitinib activity demonstrated via preclinical mouse models of asthma
 - Significant decrease of airway hyper-responsiveness
 - Significant decrease of eosinophils recruitment
- ❖ Clinical proof-of-concept in cat [Lee-Fowler, 2012] and human [Humbert, 2009] studies



Study AB07015 evaluated masitinib 6.0 mg/kg/day in severe asthma uncontrolled by OCS with no restriction on baseline eosinophil level

- Randomized (2:1), double-blinded, placebo-controlled.
- Patient with severe asthma, uncontrolled by OCS (≥ 7.5 mg/d), both high (≥ 150 cells/ μ L) and low (< 150 cells/ μ L) eosinophils
- Timing: 2-week run-in (blinded placebo) \rightarrow 36-week treatment period [W0–W36]
 \rightarrow possible blinded extension
- Primary endpoint: Reduction of annualized severe asthma exacerbation rate for overall exposure
- If significant, sequential analysis in pts with severe asthma and eosinophil count ≥ 150 cells/ μ L (fixed hierarchical procedure with 5% α for each analysis)

Study AB07015 Primary Endpoint Results



Masitinib significantly decreased the rate of severe asthma exacerbations in patients with severe asthma uncontrolled by OCS, regardless of eosinophil level

Primary Analysis (Severe Asthma)						Sequential Analysis (Severe Asthma with High Eosinophil)					
Annualized severe asthma exacerbation rate						Annualized severe asthma exacerbation rate					
	Exp	Rate	RR [95%CI]	Reduction	P-value		Exp	Rate	RR [95%CI]	Reduction	P-value
MAS (240)	1.14	0.34	0.65	35%	0.0103	MAS (181)	1.10	0.34	0.62	38%	0.0156
PBO (115)	1.15	0.48	[0.47, 0.90]			PBO (87)	1.12	0.51	[0.42, 0.91]		

Exp: Exposure in years. RR: rate ratio. MAS: Masitinib 6.0 mg/kg/d. PBO: Placebo.

- ❖ Average exposure (approx. 60 weeks) was well-balanced across treatment-arms
- ❖ Sensitivity analysis on ITT population was similarly significant (-33%, p-value=0.0156)

Sensitivity Analysis on Primary Endpoint



Benefit of masitinib was greatest in patients who had higher cumulated use of OCS (indicative of more severe asthma that is harder to control)

Sensitivity Analysis (Severe Asthma)					
Annualized severe asthma exacerbation rate					
Cumulative OCS >500 mg					
	Exp	Rate	RR [95%CI]	Reduction	P-value
MAS (161)	1.15	0.34	0.59	41%	0.0092
PBO (82)	1.20	0.55	[0.39, 0.88]		
Cumulative OCS >1000 mg					
	Rate	RR [95%CI]	Reduction	P-value	
MAS (120)	1.16	0.26	0.49	51%	0.0060
PBO (66)	1.27	0.53	[0.29, 0.82]		

Sensitivity Analysis (Severe Asthma with High Eosinophil)					
Annualized severe asthma exacerbation rate					
Cumulative OCS >500 mg					
	Exp	Rate	RR [95%CI]	Reduction	P-value
MAS (127)	1.12	0.32	0.51	49%	0.0049
PBO (60)	1.16	0.60	[0.32, 0.82]		
Cumulative OCS >1000 mg					
	Rate	RR [95%CI]	Reduction	P-value	
MAS (92)	1.11	0.22	0.29	71%	0.0003
PBO (46)	1.27	0.55	[0.15, 0.57]		

Exp: Exposure in years. RR: rate ratio. MAS: Masitinib 6.0 mg/kg/d. PBO: Placebo. OCS oral corticosteroid.



Safety was consistent with the known profile for masitinib

Summary of Adverse Events (AE) - Safety Population (n = 404)

	Masitinib % (n)	Placebo % (n)	Difference
At least one AE	83.4% (226/271)	82.0% (109/133)	+1.4%
At least one severe AE	48.0% (130/271)	45.9% (61/133)	+2.1%
At least one serious AE (non-fatal)	17.7% (48/271)	16.5% (22/133)	+1.2%

❖ No new safety signals were observed



Study AB07015 demonstrated efficacy in a difficult to treat population

❖ Study AB07015 population is distinct from other asthma trials

- Patients dependent on OCS (100% receiving high dose OCS therapy) and no weaning
- Patients were treated irrespective of baseline eosinophil count
- Evaluated over a long period of time (approx. 60 weeks)

❖ Significant reduction in severe asthma exacerbation rate

- -35% reduction in primary analysis population irrespective of baseline eosinophil level
- -38% reduction in subgroup with baseline eosinophil level ≥ 150 cells/ μL
- Greatest benefit (-41% to -71%) for patients who had the most severe asthma

❖ Masitinib may therefore provide a new treatment option for severe asthma uncontrolled by OCS

- Biologic-ineligible patients (e.g. eosinophil count of ≤ 300 cells/ μL)
- Patients in failure to biologics

Acknowledgements



THANK YOU TO OUR PATIENTS AND THEIR FAMILIES, TO ALL PARTICIPANTS AND INVESTIGATORS OF STUDY AB07015

AUTHORS: on behalf of the AB07015 STUDY GROUP

- **DAVIDESCU Lavinia: University of Oradea, Romania**
- **CHANEZ Pascal: Aix-Marseille University, France**
- **URSOL Grigoriy: Acinus, Kropyvnytskyi, Ukraine**
- **KORZH Oleksii: Kharkiv Medical Academy, Ukraine**
- **DESHMUKH Vikranth: Respira Hospital, India**
- **KURYK Lesia: Academy of Medical Science, Ukraine**
- **NORTJE Monja-Marie: Moriana Clinical Research, South Africa**
- **GODLEVSKA Olga: Kharkiv Medical Academy, Ukraine**
- **DEVOUASSOUX Gilles: Hôpital de la Croix Rousse, France.**
- **KHODOSH Eduard: City Clinical Hospital #13, Kharkiv, Ukraine.**
- **ISRAEL Elliot: Brigham and Women's Hospital, USA**
- **HERMINE Olivier: Hôpital Necker, Paris, France**



Study AB07015 was funded by AB Science, Paris, France